ry appears to be in a sheltered swamp in Manchester, Cape Ann, about thirty miles north of Boston. It here attains to but small size, and is frequently killed to the ground by severe winters. It is common in the Middle and Southern States, and Michaux informs us, that it is one of the most abundant trees in the morasses of Florida and Lower Louisiana. According to this author however, it is not usually met with far in the interior, or to the west of the mountains. Its common names are various, and change with almost every district. In Massachusetts it has no other name than *Magnolia;* in the Middle States it is called *Swamp sassafras* and *Beaver tree;* while in the Southern States it is denominated *Sweet bay* and *White bay.* It is naturally a tenant of deep boggy swamps, and is somewhat irregular in its growth. It acquires more symmetry of form when cultivated in an upland soil, although its transplantation is difficult. To insure it sussessful cultivation in a dry soil, the tree should be raised from the seed.

This tree begins to flower in different parts of the United States in May, June and July. The flowers are highly fragrant, and may be perceived by their perfume at a considerable distance. A few of them shut up in a room over night communicate to the air a heavy and almost insupportable odour.

The Magnolias are found in the class *Polyandria* and order *Polygynia;* the *Coadunatæ* of Linnæus and *Magnoliæ* of Jussieu.

This genus has a *calyx of three leaves, a corolla of six petals or more; capsules two-valved, imbricated, forming a cone; seeds berried, pendulous.*

The present species has *oval leaves, glaucous underneath;* and *obovate petals, narrowed at base.*

The bark of the young twigs is of a bright, smooth green, with rings at the insertion and scars of the leaves. The leaves are scattered, petioled, regularly elliptical, entire, and glabrous. Their under side, except the midrib, is of a beautifully pale, glaucous colour, by which the tree may be distinguished at a distance. When young, this surface is covered with a silken pubescence. Flowers solitary, terminal, on a short, incrassated peduncle. Calyx of three spatulate, obtuse, concave segments. Corolla of from eight to fourteen obovate, obtuse, concave petals, contracted at their base. The stamens are very numerous and inserted in common with the petals on the sides of a conical receptacle. Filaments very short; anthers linear, mucronated, two-celled, opening inwardly. Germs oval, collected into a cone, each one divided by a furrow and tipt with a brownish,